

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A01N 47/36 // (A01N 47/36, 43:80, 41:10, 37:42)</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/56251</b> <b>(43) International Publication Date:</b> 17 December 1998 (17.12.98)
<b>(21) International Application Number:</b> PCT/EP98/04006 <b>(22) International Filing Date:</b> 8 June 1998 (08.06.98) <b>(30) Priority Data:</b> 60/049,280 10 June 1997 (10.06.97) US <b>(71) Applicant (for all designated States except US):</b> RHONE-POULENC AGRO [FR/FR]; 14-20, rue Pierre Baizet, F-69009 Lyon (FR). <b>(72) Inventor; and</b> <b>(75) Inventor/Applicant (for US only):</b> LUFF, Kelvan, R. [US/US]; 3560 East 4000 North, Kimberly, ID 83341 (US). <b>(74) Agent:</b> BRACHOTTE, Charles, Yves; Rhône-Poulenc Agro - D.P.I., Boîte postale 9163, F-69263 Lyon Cedex 09 (FR).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> NEW HERBICIDAL MIXTURES  <b>(57) Abstract</b>  The invention relates to a method of reducing phytotoxicity to crop plants caused by at least one member of the group consisting of a 4-benzoylisoxazole herbicide and a 2-cyano-1,3-dione herbicide which comprises applying to the locus of the crop plant, the crop or crop plant seed, an antidotally effective amount of at least one sulfonyluera herbicide.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

### New Herbicidal Mixtures

#### Field of the invention.

5 This invention relates to a method of safening herbicidal  
4-benzoylisoxazoles and 2-cyano-1,3-diones by sulfonylurea herbicides, and to  
compositions containing the same.

#### Background of the invention.

10 It is known that many herbicides injure crop plants at herbicide  
application rates needed to control weed growth. This renders many herbicides  
unsuitable for controlling weeds in the presence of certain crops. Where weed  
growth in crops is uncontrolled however, this results in lower crop yield and  
reduced crop quality, as weeds will compete with crops for nutrients, light and  
water. Reduction in herbicidal injury to crops without an unacceptable  
15 reduction in the herbicidal action can be accomplished by use of crop  
protectants known as "safeners", also sometimes referred to as "antidotes" or  
"antagonists".

20 4-Benzoylisoxazoles are known to possess herbicidal properties, for  
example see European Patent Publication Nos. 0418175, 0487357, 0527036 and  
0560482. European Patent Publication Nos. 0496630, 0496631, 0625505 and  
0625508 disclose certain 1-phenyl-2-cyano-1,3-dione derivatives possessing  
herbicidal properties. European Patent Publication No. 0213892 discloses  
herbicidally active enols. These compounds possess very good levels of  
herbicidal activity, but at higher dose rates there can be a risk of crop  
25 phytotoxicity.

The present invention seeks to provide compositions of these herbicides  
for use in combination with antidotes therefor for reduction of injury to crops,  
especially wheat, due to phytotoxicity of these herbicides at certain dose rates or  
under certain conditions of use.

- 2 -


Description of the Invention

The invention provides a method of reducing phytotoxicity at a crop plant locus caused by at least one 4-benzoylisoxazole herbicide and/or 2-cyano-1,3-dione herbicide which comprises applying to the crop plant locus at least one sulfonylurea herbicide.

Suprisingly, the applicants have found that the presence of at least one sulfonylurea herbicide allows any crop phytotoxicity by the isoxazole and/or 2-cyano-1,3-dione herbicide to be reduced in the presence of at least one sulfonylurea herbicide, while maintaining a good level of weed control.

The method of the invention reduces phytotoxicity by a safening effect of the sulfonylurea. The invention also provides a method whereby the total amount of herbicide may be reduced by virtue of a synergistic effect.

Preferably the 4-benzoylisoxazole herbicide has the general formula (I):



(I)

wherein

R is hydrogen or  $-\text{CO}_2\text{R}^3$ ;

$\text{R}^1$  is  $\text{C}_{1-6}$  alkyl or  $\text{C}_{3-6}$  cycloalkyl optionally bearing  $\text{C}_{1-6}$  alkyl;

$\text{R}^2$  is selected from halogen (e.g. chlorine or bromine),  $-\text{S}(\text{O})_p\text{Me}$

$\text{CH}_2\text{SO}_q\text{Me}$ ,  $\text{C}_{1-6}$  alkyl,  $\text{C}_{1-6}$  haloalkyl (e.g.  $\text{CF}_3$ ),  $\text{C}_{1-6}$  alkoxy or  $\text{C}_{1-6}$  haloalkoxy;

n is two or three; p is zero, one or two; q is zero, one or two; and

$\text{R}^3$  is  $\text{C}_{1-4}$  alkyl.

In formula (I) above  $\text{R}^1$  is preferably cyclopropyl.

In formula (I) above preferably one group  $\text{R}^2$  represents  $-\text{S}(\text{O})_p\text{Me}$ , most preferably in the 2-position of the benzoyl ring.

- 3 -


4-Benzoylisoxazoles of formula (I) above of particular interest in the method of the invention include the following:

- A. 5-cyclopropyl-4-(2-methylsulfonyl-4-trifluoromethyl)benzoylisoxazole;
- 5 B. ethyl 5-cyclopropyl-4-(2-methylsulfonyl-4-trifluoromethyl)benzoylisoxazole-3-carboxylate;
- C. ethyl 5-cyclopropyl-4-[3,4-dichloro-2-(methylsulphenyl)benzoyl]isoxazole-3-carboxylate;
- D. 5-cyclopropyl-4-[4-bromo-2-(methylsulfonylmethyl)benzoyl]isoxazole;
- 10 E. 5-cyclopropyl-4-(4-methylsulfonyl-2-trifluoromethyl)benzoylisoxazole; and
- F. 5-cyclopropyl-4-(4-chloro-2-methylsulfonyl)benzoylisoxazole.

The letters A to F are assigned to these compounds for reference and identification hereafter.

Compounds A, B and C are preferred.

Preferably the 2-cyano-1,3-dione derivative has the formula (II):



(II)

20 wherein  $R^1$ ,  $R^2$  and  $n$  are as defined above.

Compounds of formula II above may exist in enolic tautomeric forms that may give rise to geometric isomers around the enolic double bond.

Furthermore, in certain cases the groups  $R^1$  and  $R^2$  may give rise to stereoisomers and geometric isomers. All such forms and mixtures thereof are embraced by the present invention.

25

- 4 -

The most preferred compound of formula (II) above is 2-cyano-3-cyclopropyl-1-(2-methylsulfonyl-4-trifluoromethylphenyl)propan-1,3-dione.

Preferably the sulfonylurea herbicide has the general formula (III):



5            wherein:

$R_4$  is selected from substituted or unsubstituted phenyl, substituted or unsubstituted thienyl, substituted or unsubstituted pyridyl, and substituted or unsubstituted imidazopyridinyl;

$R_5$  and  $R_6$  are independently  $C_{1-6}$  alkyl or hydrogen; or

10             $R_7$  is selected from substituted or unsubstituted triazine or substituted or unsubstituted diazine. Preferably the substituted phenyl, thienyl, pyridyl or imidazopyridinyl moieties are substituted by halogen, alkyl, alkoxy, alkylcarbonyl, alkoxycarbonyl, substituted or unsubstituted amido, alkylthio, alkylsulfenyl, and alkylsulfonyl. The triazine and diazine moieties are preferably  
15            substituted by alkoxy or alkyl.

In the above definition, alkyl and moieties comprising it generally contain from one to six carbon atoms and are optionally substituted by one or more atoms from group consisting of bromine, chlorine, fluorine and iodine.

20            Preferably the sulfonylurea herbicide has the general formula (III) with one or more of the following features:

$R_4$  is selected from 2-substituted phenyl, 2-substituted-3-thienyl, 3-substituted-2-pyridyl; 2-substituted-imidazo[1,2-a]pyridine;

$R_5$  and  $R_6$  are independently methyl or hydrogen; and

25             $R_7$  is selected from 4,6-dimethoxy-1,3,5-triazine; 4-methoxy-6-methyl-1,3,5-triazine; 4,6-bisdifluoromethoxy-1,3-diazine, and 4,6-dimethoxy-1,3-diazine.

Even more preferably the sulfonylurea herbicide has the general formula (III) wherein:

- 5 -

R<sub>4</sub> is selected from 2-chlorophenyl; 2-methoxycarbonylphenyl, 2-haloalkylphenyl, 2-haloalkoxyphenyl, 2-methoxycarbonyl-3-thienyl, 3-dialkylamido-2-pyridyl and 2-alkylsulfonylimidazo[1,2-a]pyridine;

R<sub>5</sub> and R<sub>6</sub> are independently selected from hydrogen or methyl; and

5 R<sub>7</sub> is selected from 4,6-dimethoxy-1,3,5-triazine; 4-methoxy-6-methyl-1,3,5-triazine; 4,6-bisdifluoromethoxy-1,3-diazine; and 4,6-dimethoxy-1,3-diazine.

Preferably there are one or two sulfonylurea herbicides present selected from:

10 1-(2-chlorophenylsulfonyl)-3-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)urea (chlorsulfuron);

2-[4-methoxy-6-methyl-1,3,5-triazin-2-yl(methyl)carbamoylsulfamoyl]benzoic acid (tribenuron-methyl);

15 3-(4-methoxy-6-methyl-1,3,5-triazin-2-ylcarbamoylsulfamoyl)thiophen-2-carboxylic acid (thifensulfuron methyl);

2-(4-methoxy-6-methyl-1,3,5-triazin-2-ylcarbamoylsulfamoyl)benzoic acid (metsulfuron);

1-[2-(2-chloroethoxy)phenylsulfonyl]-3-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)urea (triasulfuron);

20 1-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-3-[2-(3,3,3-trifluoropropyl)-phenylsulfonyl]urea (CGA-152005 or prosulfuron);

1-(4,6-dimethoxypyrimidin-2-yl)-3-(3-dimethylcarbamoyl-2-pyridylsulfonyl)urea (nicosulfuron);

25 2-[4,6-bis(difluoromethoxy)pyrimidin-2-yl-carbamoylsulfamoyl]benzoic acid (primisulfuron);

N-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]-3-ethylsulfonyl-2-pyridinesulfonamide (rimsulfuron or DPX-E9636) and

- 6 -

1-(2-ethylsulfonylimidazo[1,2-a]pyridin-3-ylsulfonyl)-3-(4,6-dimethoxypyrimidin-2-yl)urea (MON 37500: CAS Registry Number 141776-32-1).

5       Sulfonylurea herbicides may be prepared by references found in The Pesticide Manual, 10<sup>th</sup> Edition, C. Tomlin editor; British Crop Protection Association, 1994, or by those known to the skilled addressee.

      Most preferably there are two sulfonyl urea herbicides present. Preferably these are selected from thifensulfuron methyl, tribenuron methyl and MON 37500.

10       The amount of sulfonylurea herbicide used in the method of the invention varies according to a number of parameters including the weeds to be controlled, the crop to be protected, the amount and rate of herbicide applied, and the edaphic and climatic conditions prevailing. Also, the selection of the specific antidotes for use in the method of the invention, the manner in which it is to be applied and the determination of the activity which is non-phytotoxic but antidotally effective, can be readily performed in accordance with common practice in the art.

15       By "non-phytotoxic" is meant an amount of the antidote which causes at most minor or no injury to the desired crop species. By "antidotally effective" is meant an antidote used in an amount which is effective as an antidote to decrease the extent of injury caused by the herbicide to the desired crop species.

20       The dose rate of the benzoylisoxazole herbicide and/or 2-cyano-1,3-dione herbicide is generally from about 5 to about 500 grammes per hectare (g/ha), preferably from about 15 to about 200 g/ha, more preferably from about 20 to about 120 g/ha, even more preferably from about 70 to about 90 g/ha.

25       The dose rate of the sulfonylurea herbicide is generally from about 1 to about 250 g/ha, preferably from about 1 to about 100 g/ha, more preferably from about 3 to about 20 g/ha and even more preferably from about 5 to about 15 g/ha.



- 7 -

Preferably the weight ratio of isoxazole/dione:sulfonyurea herbicide is from about 500:1 to about 1:50, preferably from about 200:1 to about 1:7, more preferably from about 40:1 to about 1:1, and even more preferably from about 18:1 to about 5:1.

5           The method of the invention can be applied pre- or post-emergence of the crop. Where the crop is a cereal crop (such as wheat) the herbicide is preferably applied post-emergence of the crop species.

According to a further feature of the present invention there is provided a composition comprising:

10           (a) at least one 4-benzoylisoxazole herbicide or 2-cyano-1,3-dione herbicide; and

(b) at least one sulfonylurea herbicide,

15           in association with an agriculturally acceptable diluent or carrier and/or optionally a surface active agent. The composition may be provided as a ready-to-use formulation (e.g. where (a) and (b) are premixed), or may be formed as a tank mix in accordance with standard techniques in the art.

The herbicidal action of the composition may be substantially higher than the sum of the effects of the individual agents. The effect may be a synergistic effect.

20           The composition according to the invention can be used both in conventional methods of cultivation (strip cultivation with suitable strip width) and in plantation cultivation (e.g., vines, fruit, citrus), as well as in industrial plants and track systems, on roads and squares, but also to handle stubble and in the minimum-tillage method. They are also suitable as burners (for killing  
25           foliage, e.g. in potatoes) or as defoliants (e.g. in cotton). They are also suitable for use on fallow areas. Other areas of use are in tree nurseries, forests, grasslands, and in the cultivation of ornamental plants.

Examples of weeds which can be combatted well by the agents or combinations of agents according to the invention are:

- 8 -

Dicotyledon weeds of the genera sinapis, lepidium, galium, stellaria, matricaria, anthemis, galinsoga, chenopodium, urtica, senecio, amaranthus, portulaca, xanthium, convolvulus, ipomoea, polygonum, sesbania, ambrosia, solanum, cirsium, carduus, sonchus, rorippa, rotaia, lindernia, lamium,  
5 veronica, abutilon, emex, sida, datura, viola, galeopsis, papaver, centaurea, trifolium, ranunculus, taraxum, and mentha.

Monocotyledon weeds of the genera echinocloa, setaria, panicum, digitaria, phleum, poa, festuca, eleusine, brachiaria, lolium, bromus, avena, cyperus, sorghum, agropyron, cynodon, monochoria, fimbristylis, sagittaria,  
10 eleocharis, scirpus, papalum, ischaemum, spencoclea, dactyloctenium, agrostis, alopecurus, apera.

However, the use of the agents and combinations of agents according to the invention is in no way limited to these genera, but rather extends in the same way to other plants.

15 The crops that may be protected by the method of the invention include corn, rice, wheat, soya, sorghum and cotton. The method of the invention is preferably performed where the crop to be protected is wheat.

The invention also provides a product comprising at least one 4-benzoylisoxazole herbicide and/or 2-cyano-1,3-dione herbicide and a  
20 sulfonylurea herbicide for simultaneous, separate or sequential use in the control of weeds at a locus.

The following non-limiting Examples illustrate the invention.

### Example 1

25 Compositions containing Compound A (formulated as a wettable powder containing 75 % active ingredient) alone, in tank-mixture with Triton Ag 98 (trademark, 0.25% v/v; a non-ionic surfactant) and in tank-mixture with Triton Ag 98, thifensulfuron methyl and tribenuron methyl were mixed together in a spray tank (the latter two sulfonylurea herbicides were used as the commercial

- 9 -

formulation HARMONY EXTRA™ a wettable powder containing 75% combined active ingredients) were sprayed at a volume of 225 liters per hectare over Penawawa Soft White Spring Wheat which had emerged from seeding and grown to a height of about 18 cm. Comparative ratings for damage (by visual inspection of the degree of chlorosis present) to the wheat were taken at 4 and 8 days after treatment (DAT) in comparison with untreated control. The following results were observed:

Mixture	Rate Active Ingredient (g/ha)	Percent Chlorosis	
		4 DAT	8 DAT
Compound A	70	0	0
Compound A+ Triton AG 98	70	23	60
Compound A + Triton AG 98 thifensulfuron methyl tribenuron methyl	70 11.6 5.8	15	37

### Example 2

The same procedure as described in Example 1 was followed replacing Compound A with Compound B (which was formulated as a suspension concentrate containing 4% active ingredient). The following results were obtained.

Mixture	Rate Active Ingredient (g/ha)	Percent Chlorosis	
		4 DAT	8 DAT
Compound B	85	0	0
Compound B + Triton Ag 98	85	18	38
Compound B + Triton Ag 98 thifensulfuron methyl tribenuron methyl	85 11.6 5.8	11	20

- 10 -

**Example 3**

Compound C as a 25% emulsifiable concentrate and prosulfuron (as a 20% wettable powder) were suspended in tap water and applied post-emergence both alone and in combination at a range of concentrations to Xanthium strumarium,  
5 Amaranthus retroflexus, and Setaria faberi. Treatment effects were assessed visually 20 days after treatment. The percentage damage compared to untreated controls was recorded for each species.

The nature of the interaction between the two components was determined using the responses of the herbicides applied singly in calculating the expected  
10 response when they are combined (COLBY, S.R., "Calculating synergistic and antagonistic response of herbicide combinations". Weeds 15, pages 20-22, 1967):

$$We = X + \frac{Y \cdot (100 - X)}{100}$$

15 wherein

X = Percentage mortality, compared with untreated controls, after treatment with Compound C at a rate of application of p g/hectare.

20 Y = Percentage mortality, compared with untreated controls, after treatment with prosulfuron at a rate of application of q g/hectare.

We = The expected pesticidal activity (percentage mortality compared with untreated controls) after treatment with Compound C and prosulfuron at a rate of  
25 application of p + q g/ha.

In the tables that follow the figures used for weed control are percentages reduction in growth when compared with untreated controls. The figures in parentheses are those expected using the Colby formula.

30

- 11 -

**Table 1**

**Post-emergence treatment of Xanthium strumarium with various mixtures of Compound C and prosulfuron**

5

Prosulfuron

Cpd C	Dose g/ha	0	1	2	4	8
	0	-	0	15	0	55
	8	0	5 ( 0 )	15 ( 15 )	25 ( 0 )	65 ( 55 )
	16	5	25 ( 5 )	40 ( 19 )	60 ( 5 )	90 ( 57 )
	32	15	20 ( 15 )	45 ( 28 )	45 ( 15 )	87 ( 62 )
	63	20	30 ( 20 )	35 ( 32 )	62 ( 20 )	77 ( 64 )
	125	40	55 ( 40 )	85 ( 49 )	80 ( 40 )	95 ( 73 )

10

**Table 2**

**Post-emergence treatment of Amaranthus retroflexus with various mixtures of Compound C and prosulfuron**

15

Prosulfuron

Cpd C	Dose g/ha	0	4	8
	0	-	20	35
	8	20	60 ( 36 )	77 ( 48 )
	16	40	60 ( 52 )	75 ( 61 )
	32	52	82 ( 62 )	87 ( 69 )

- 12 -

**Table 3**

**Post-emergence treatment of Setaria faberi with various mixtures of Compound C and prosulfuron**

5

Prosulfuron

Cpd C	Dose g/ha	0	1	2	4	8	16
	0	-	0	5	20	20	32
	16	35	50 ( 35 )	50 ( 38 )	60 ( 48 )	60 ( 48 )	75 ( 56 )

10 For a range of mixtures of Compound C with prosulfuron against Xanthium strumarium, Amaranthus retroflexus and Setaria faberi the observed response was greater than expected response thus indicating synergism.

15 While the invention has been described in terms of various preferred embodiments, the skilled artisan will appreciate that various modifications, substitutions, omissions, and changes may be made without departing from the spirit thereof. Accordingly, it is intended that the scope of the present invention be limited solely by the scope of the following claims, including equivalents thereof.

20

- 13 -


CLAIMS

1. A method of reducing phytotoxicity at a crop plant locus caused by at least one 4-benzoylisoxazole herbicide and/or 2-cyano-1,3-dione herbicide which comprises applying to the crop plant locus at least one sulfonylurea herbicide.

2. The method according to Claim 1 wherein phytotoxicity is reduced by a safening effect of the sulfonylurea.

3. The method according to Claim 1 or Claim 2 wherein the total amount of herbicide is reduced by virtue of a synergistic effect.

4. The method according to any one of the foregoing claims wherein the 4-benzoylisoxazole has the general formula (I):



(I)

wherein

R is hydrogen or  $-\text{CO}_2\text{R}^3$ ;

$\text{R}^1$  is  $\text{C}_{1-6}$  alkyl or  $\text{C}_{3-6}$  cycloalkyl optionally bearing  $\text{C}_{1-6}$  alkyl;

$\text{R}^2$  is selected from halogen,  $-\text{S}(\text{O})_p\text{Me}$ ,  $\text{CH}_2\text{S}(\text{O})_q\text{Me}$ ,  $\text{C}_{1-6}$  alkyl,  $\text{C}_{1-6}$  haloalkyl,  $\text{C}_{1-6}$  alkoxy and  $\text{C}_{1-6}$  haloalkoxy;


n is two or three; p is zero, one or two; q is zero, one or two; and

$\text{R}^3$  is  $\text{C}_{1-4}$  alkyl.

- 14 -

5. The method according to any one of the foregoing claims in which the 4-benzoylisoxazole is 5-cyclopropyl-4-(2-methylsulfonyl-4-trifluoromethyl)benzoylisoxazole.

6. The method according to any one of Claims 1 to 3 in which the 2-cyano-1,3-dione herbicide has the formula (II):



(II)

wherein

R<sup>1</sup> is C<sub>1-6</sub> alkyl or C<sub>3-6</sub> cycloalkyl optionally bearing C<sub>1-6</sub> alkyl;

R<sup>2</sup> is selected from halogen, -S(O)<sub>p</sub>Me and C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy and C<sub>1-6</sub> haloalkoxy;

n is two or three; and p is zero, one or two.

7. The method according to any one of the foregoing claims in which the crop plant to be protected is corn, rice, wheat, soya, sorghum, and cotton; preferably wheat.

8. The method according to any one of the foregoing claims in which the dose rate of the benzoylisoxazole herbicide and/or 2-cyano-1,3-dione herbicide is from about 5 to about 500 g/ha.

9. The method according to any one of the foregoing claims wherein the sulfonylurea herbicide has the general formula (III):



wherein:



- 15 -

R<sub>4</sub> is selected from substituted or unsubstituted phenyl, substituted or unsubstituted thienyl, substituted or unsubstituted pyridyl, and substituted or unsubstituted imidazopyridine;

R<sub>5</sub> and R<sub>6</sub> are independently C<sub>1-6</sub> alkyl or hydrogen; and

5 R<sub>7</sub> is selected from substituted or unsubstituted triazine or substituted or unsubstituted diazine.

10. The method according to Claim 9 wherein the sulfonylurea is:

10 1-(2-chlorophenylsulfonyl)-3-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)urea;

2-[4-methoxy-6-methyl-1,3,5-triazin-2-yl(methyl)carbamoylsulfamoyl]benzoic acid;

3-(4-methoxy-6-methyl-1,3,5-triazin-2-ylcarbamoylsulfamoyl)thiophen-2-carboxylic acid;

15 2-(4-methoxy-6-methyl-1,3,5-triazin-2-ylcarbamoylsulfamoyl)benzoic acid;

1-[2-(2-chloroethoxy)phenylsulfonyl]-3-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)urea;

20 1-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-3-[2-(3,3,3-trifluoropropyl)-phenylsulfonyl]urea;

1-(4,6-dimethoxypyrimidin-2-yl)-3-(3-dimethylcarbamoyl-2-pyridylsulfonyl)urea;

2-[4,6-bis(difluoromethoxy)pyrimidin-2-yl-carbamoylsulfamoyl]benzoic acid;

25 N-[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]-3-ethylsulfonyl-2-pyridinesulfonamide; or

1-(2-ethylsulfonylimidazo[1,2-a]pyridin-3-ylsulfonyl)-3-(4,6-dimethoxypyrimidin-2-yl)urea.

- 16 -

11. A composition comprising:

(a) at least one 4-benzoylisoxazole herbicide or 2-cyano-1,3-dione herbicide; and

5 (b) at least one sulfonylurea herbicide in association with an agriculturally acceptable diluent or carrier.

12. The composition according to Claim 11 in which the 4-benzoylisoxazole is 5-cyclopropyl-4-(2-methylsulfonyl)-4-trifluoromethyl)benzoylisoaxole.

10

13. A product comprising at least one 4-benzoylisoxazole herbicide and/or 2-cyano-1,3-dione herbicide and a sulfonylurea herbicide for simultaneous, separate or sequential use in the control of weeds at a locus.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 98/04006

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A01N47/36 //(A01N47/36, 43:80, 41:10, 37:42)

According to International Patent Classification(IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
------------	--	-----------------------

A	WO 96 21357 A (ZENECA LTD) 18 July 1996 -----	
---	--	--

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

21 October 1998

Date of mailing of the international search report

29/10/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Decorte, D

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/04006

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9621357 A	18-07-1996	US 5627131 A	06-05-1997
		AU 4312996 A	31-07-1996
		CA 2209937 A	18-07-1996
		CN 1168083 A	17-12-1997
		CZ 9702068 A	13-05-1998
		EP 0802732 A	29-10-1997
		HU 9800117 A	28-04-1998
		JP 10505099 T	19-05-1998
		NZ 297874 A	26-08-1998
		PL 321266 A	24-11-1997
		ZA 9600067 A	18-07-1996